

RESEARCH ARTICLE

Open Access



Photoinactivation of major bacterial pathogens in aquaculture

Heyong Jin Roh, Ahran Kim, Gyoung Sik Kang and Do-Hyung Kim*

Abstract

Background: Significant increases in the bacterial resistance to various antibiotics have been found in fish farms. Non-antibiotic therapies for infectious diseases in aquaculture are needed. In recent years, light-emitting diode technology has been applied to the inactivation of pathogens, especially those affecting humans. The purpose of this study was to assess the effect of blue light (wavelengths 405 and 465 nm) on seven major bacterial pathogens that affect fish and shellfish important in aquaculture.

Results: We successfully demonstrate inactivation activity of a 405/465-nm LED on selected bacterial pathogens. Although some bacteria were not fully inactivated by the 465-nm light, the 405-nm light had a bactericidal effect against all seven pathogens, indicating that blue light can be effective without the addition of a photosensitizer. *Photobacterium damsela*, *Vibrio anguillarum*, and *Edwardsiella tarda* were the most susceptible to the 405-nm light (36.1, 41.2, and 68.4 J cm⁻², respectively, produced one log reduction in the bacterial populations), whereas *Streptococcus parauberis* was the least susceptible (153.8 J cm⁻² per one log reduction). In general, optical density (OD) values indicated that higher bacterial densities were associated with lower inactivating efficacy, with the exception of *P. damsela* and *Vibrio harveyi*. In conclusion, growth of the bacterial fish and shellfish pathogens evaluated in this study was inactivated by exposure to either the 405- or 465-nm light. In addition, inactivation was dependent on exposure time.

Conclusions: This study presents that blue LED has potentially alternative therapy for treating fish and shellfish bacterial pathogens. It has great advantages in aspect of eco-friendly treating methods differed from antimicrobial methods.

Keywords: Photoinactivation, Blue light, Bacterial fish pathogen, Fish disease, Light-emitting diode

Abbreviations: BE, Bactericidal efficiency; BHIA, Brain and heart infusion agar; BHIB, Brain and heart infusion broth; LED, Light-emitting diode; OD, Optical density; PBS, Phosphate buffered saline; PPF, Photosynthesis photon flux density; TSA, Tryptic soy agar

Background

Aquaculture has been the fastest-growing food-producing sector since 1970, with an average growth rate of ~9 % per year, compared with a 2.8 % growth rate of terrestrial farmed meat production over the same period (Bostock et al. 2010; Subasinghe et al. 2001). Worldwide, disease is considered to be a significant constraint on aquaculture; the economic losses caused by disease are estimated to be several billion US dollars per year (Subasinghe et al. 2001). Bacterial diseases are a major threat to aquaculture

because bacteria can survive well and reach high densities in an aquatic environment independent of their hosts, which is generally not the case in terrestrial environments (Defoirdt et al. 2011; Pridgeon and Klesius 2013). In particular, the larval stages of several farmed aquatic animals are highly susceptible to bacterial diseases (Defoirdt et al. 2011). Major bacterial pathogens include *Vibrio*, *Aeromonas*, *Edwardsiella*, and *Streptococcus* species, which affect fish such as salmon, carp, and flat fish (Baeck et al. 2006; Han et al. 2006; Milton et al. 1996; Romalde 2002; Weinstein et al. 1997; Wiklund and Dalsgaard 1998; Won and Park 2008). Inactivation of microorganisms can be accomplished with light

* Correspondence: dhkim@pknu.ac.kr
Department of Aquatic life Medicine, College of Fisheries Science, Pukyong National University, Busan, South Korea

technologies, including ultraviolet C irradiation therapy, photodynamic therapy (PDT), and blue light therapy (Arrojado et al. 2011; Yin et al. 2013). Ultraviolet (UV) irradiation has an adverse effect on fish; it causes intensive skin lesions (Ghanizadeh and Khodabandeh 2010) and reduction of goblet cells in fish skin, resulting in less mucus production and, consequently, downregulation of innate immunity (Kaweewat and Hofer 1997). The use of blue light (400–500 nm) as a mono-therapy is gaining increasing attention because of its potential antimicrobial effect and because it does not require an exogenous photosensitizer (Yin et al. 2013). Blue light is much less harmful to mammalian cells than UV irradiation (Kleinpenning et al. 2010). Light treatment has been applied in aquaculture for many years. For example, European sea bass and sole larvae showed the fastest development and the lowest degree of deformity under blue light (half-peak bandwidth = 435–500 nm) than under other wavelengths of light (Villamizar et al. 2011). Also, another study found that retina from fish exposed to blue light revealed no signs of damage as assessed by extensive histological examination (Migaud et al. 2007). In spite of this potential, there is little information on light therapy as it applies to bacterial pathogens that threaten aquaculture. The aim of this study was to determine the extent of inactivation of bacterial fish pathogens, in particular, seven species including both Gram-negative and Gram-positive bacteria carried out in *in vitro* experiment. The effects of light-emitting diode (LED) on different bacterial densities and the effects of different light intensities were also evaluated.

Methods

Bacterial strains and identification

Seven bacterial species were evaluated in this study. The bacterial strains were grown on tryptic soy agar (TSA) or brain and heart infusion agar (BHIA), supplemented with 1 % NaCl. A strain of *Vibrio anguillarum* isolated from diseased cod was purchased from the Korean collection for type cultures (KCTC), and *Edwardsiella tarda* KE1 and *Aeromonas salmonicida* RFAS1 originated from diseased olive flounder and black rockfish were previously used (Han et al. 2006, 2011). *Vibrio harveyi* Vh21FL, *Photobacterium damsela* Dae1-1L, *Streptococcus iniae* BS9, and *Streptococcus parauberis* SpOF3K obtained from diseased olive flounder were confirmed by polymerase chain reaction that was previously described (Table 1) (Mata et al. 2004; Osorio et al. 2000; Pang et al. 2006).

LED source

The 405- and 465-nm LEDs, each composed of 120 individual LEDs, were kindly provided by the LED-Marine Convergence Technology R&D Center (Pukyong National University). The spectra of the 405- and 465-nm LEDs as

measured by a temperature-controllable integrating system (Withlight Co. Ltd., Korea) are shown in Fig. 1. The maximum irradiation of the 405- and 465-nm LED array were 250 and 516 $\mu\text{mol m}^{-2} \text{s}^{-1}$, respectively, as calculated using a laboratory radiometer (Biospherical Instruments Inc., USA). Photosynthesis photon flux density (PPFD; $\mu\text{mol m}^{-2} \text{s}^{-1}$) was converted to radiant flux density (mW cm^{-2}) by using the following formula:

$$\text{Radiant flux}(W) = h \times C \times NA \times \text{PPFD}(\mu\text{mol})/\lambda \times 10^{-3}$$

$$h(\text{Plank's constant}) = 6.626 \times 10^{-34}$$

$$C(\text{Light velocity}) = 3 \times 10^8 \text{ms}^{-1} \quad \lambda = \text{Wavelength}(\text{nm})$$

$$NA(\text{Avogadro's constant}) = 6.02 \times 10^{23}$$

Antibacterial activity of LEDs

Approximately 10^5 CFU ml^{-1} of each culture was suspended in phosphate buffered saline (PBS; pH 7.2–7.4). Each bacterial suspension (10 ml, with a depth of 5 mm) was plated on a 30-mm petri dish on TSA (*V. anguillarum*, *V. harveyi*, *P. damsela*, *E. tarda*, and *A. salmonicida*) or BHIA (*S. iniae* and *S. parauberis*) supplemented with 1 % NaCl, exposed to 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of the 405- or 516 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of the 465-nm LED light, and placed in a 25 °C incubator for 0, 1, 3, 6, 12, 24, or 48 h. Each lamp was placed 3.5 cm above open plates containing the bacterial cultures and positioned perpendicularly. Temperature was routinely monitored during irradiation. The cultures were stirred with a sterile magnetic bar for a few seconds just before being plated, and bacterial counts were performed. A method slightly modified from a previous study (Maclean et al. 2009) was used to express the inactivation data: $\log_{10}(N/N_0)$ was plotted as a function of exposure time, where N_0 is the initial bacterial population in CFU ml^{-1} prior to inactivation and N is 10 CFU ml^{-1} . Thus, the mean bactericidal efficiency (BE) was defined as the \log_{10} reduction in a bacterial population [$\log_{10}(10/N_0)$] by inactivation per unit dose in J cm^{-2} . Exposure time was deduced from the time at which bacterial populations reached 10 CFU ml^{-1} .

In order to determine the effects of initial bacterial density on the antibacterial activity of LEDs, 200 μl of six 10-fold serial dilutions (10^3 , 10^4 , 10^5 , 10^6 , 10^7 , and 10^8 CFU ml^{-1} , in BHIB supplemented with 1 % NaCl) were inoculated in a 96-well microplate. The plates were exposed to a 405- or 465-nm LED at 25 °C. Optical density (OD) was measured at 630 nm after 24 h irradiation using a Sunrise™ spectrophotometer (TECAN Austria), and data was analyzed using OD of 24 h exposure group/OD of 24 h non-exposure group $\times 100$ (%) formula.

The data points shown in Fig. 2 and in Table 3 are expressed as mean values with standard deviations. Two-tailed Student's *t* tests and ANOVA Tukey's test were used

Table 1 Bacterial strains and primers used in this study

Bacterial strains	Isolation source	Primers	References
Gram-negative bacteria			
<i>Vibrio harveyi</i> Vh21FL	Diseased olive flounder	Tox F: 5'-GAAGCAGCACTCACCGAT-3' Tox R: 5'-GGTGAAGACTCATCAGCA-3'	Pang et al. (2006)
<i>Vibrio anguillarum</i> KCTC 2711 ^a	Diseased cod	–	–
<i>Photobacterium damsela</i> Dae1-1L	Diseased olive flounder	Car 1: 5'-GCTTGAAGAGATTGAGT-3' Car 2: 5'-CACCTCGCGGTCTTGCTG-3' Ure -3: 5'-CTTGAATATCCATCTCATCTGC-3' Ure -5: 5'-TCCGGAATAGGTAAGCGGG-3'	Osorio et al. (2000)
<i>Edwardsiella tarda</i> KE1	Diseased olive flounder	–	Han et al. (2006)
<i>Aeromonas salmonicida</i> RFAS1	Disease black Rockfish	–	Han et al. (2011)
Gram-positive bacteria			
<i>Streptococcus iniae</i> BS9	Diseased olive flounder	Lox-1: 5'-AAGGGGAAATCGCAAGTGCC-3' Lox-2: 5'-ATATCTGATTGGCCGTCTAA-3'	Mata et al. (2004)
<i>Streptococcus parauberis</i> SpOF3K	Diseased olive flounder	Spa F: 5'-TTTCGTCTGAGGCAATGTTG-3' Spa R: 5'-GCTTCATATATCGCTATACT-3'	

^aType strain

to determine statistically significant differences ($P < 0.05$ or $P < 0.01$) between groups exposed to blue light and controls.

Results

This study successfully demonstrates the bactericidal effects of 405- and 465-nm LEDs on selected bacterial fish and shellfish pathogens. As shown in Fig. 2, growth of the pathogens evaluated was clearly inactivated by exposure to either a 405- or 465-nm LED, although the degree of inactivation varied depending on bacterial species and sampling time point. The one exception was that a 465-nm LED was unable to inactivate *V. harveyi*, but that strain was inactivated by a 405-nm LED. Complete inactivation of *A. salmonicida* and *S. parauberis* was seen 24 h after irradiation with a 405-nm LED, whereas only 6 h were required for complete inactivation of *V. anguillarum* and *P. damsela* under the same conditions. Although *S. iniae* was more rapidly inhibited by a 465-nm LED, overall, there were no differences between 405 and 465 nm LEDs in the inactivation rate of *S. parauberis*.

BE was measured in this study using a method modified from one that was previously described (Maclean et al. 2009). Details of the inactivation parameters for all bacterial species are listed in decreasing order of BE in Table 2. We calculated BE using exposure time, which was deduced from the time at which bacterial populations reached 10 CFU ml⁻¹. *P. damsela*, *V. anguillarum*, and *E. tarda* were the most susceptible bacteria, while *S. parauberis* was the least susceptible, to exposure to a 405-nm LED. Our results show that Gram-negative bacteria, such as *P. damsela* (36.1 J cm⁻²), *V. anguillarum* (41.2 J cm⁻²), and *E. tarda* (68.4 J cm⁻²), seem to be more sensitive to a 405-nm LED light than are Gram-positive bacteria like *S. parauberis* (153.8 J cm⁻²) and *S. iniae* (90.4 J cm⁻²) (Table 2). However, some Gram-negative bacteria such as *A. salmonicida* (98.7 J cm⁻²) and *V. harveyi* (126.4 J cm⁻²) have lower susceptibility than *S. iniae*.

The degree of inactivation of bacterial suspensions with varying initial population densities in BHIB + 1 % NaCl following exposure to a 405- or 465-nm LED for




Fig. 1 Emission spectra of 405- (a) and 465-nm (b) LED arrays




Fig. 2 Viable bacterial counts of *V. anguillarum* (a), *V. harveyi* (b), *P. damsela* (c), *E. tarda* (d), *A. salmonicida* (e), *S. parauberis* (f), and *S. iniae* (g) in phosphate buffered saline at 25 °C and several sampling time points (1, 3, 6, 12, 24, and 48 h) after LED exposure in a 405- or 465-nm LED (respectively, 250 μ mol m⁻² s⁻¹ or 516 μ mol m⁻² s⁻¹) * significant difference, P<0.05; **significant difference, P<0.01

24 h is displayed in Table 3. In general, the OD values indicate that the higher starting bacterial densities were associated with lower inactivating efficacies. However, there were exceptions: unlike the other bacterial species, *P. damsela* exposed to a 405- or 465-nm LED and *V. harveyi* exposed to a 465-nm LED were not affected by their initial concentrations. *P. damsela* was able to survive a 405- or 465-nm light exposure in BHIB + 1 % NaCl, but it was much more susceptible when suspended in PBS.

Discussion

Antimicrobials are commonly used in aquaculture to prevent and treat bacterial infections in fish. Significant increases in the bacterial resistance to various antibiotics, such as oxytetracycline, quinolones, and amoxicillin, have been repeatedly found in proximity to fish farms (Defoirdt et al. 2011; Guardabassi et al. 2000; Schmidt et al. 2000). Excessive use of antimicrobials may significantly reduce their effectiveness and their usefulness in aquaculture. More importantly, studies have

Table 2 Energy levels and bactericidal efficiencies for the inactivation of bacterial species using 405- and 465-nm LEDs

Species	Wavelength (nm)	Dose (J cm ⁻²)	Log ₁₀ reduction (±SD)	Dose/log ₁₀ reduction (±SD)	BE ^a (±SD)
<i>Photobacterium damsela</i>	405	137 (±4)	3.8	36.1 (±1.0)	0.028 (±0.001)
	465	1387 (±126)	3.7	374.7 (±34.1)	0.003 (±0)
<i>Vibrio anguillarum</i>	405	132 (±4)	3.2	41.2 (±1.3)	0.024 (±0.001)
	465	1934 (±3)	4	483.6 (±0.7)	0.002 (±0)
<i>Edwardsiella tarda</i>	405	260 (±10)	3.8	68.4 (±2.7)	0.015 (±0.001)
	465	2178 (±517)	4.0	544.4 (±129.2)	0.002 (±0)
<i>Streptococcus iniae</i>	405	262 (±5)	2.9	90.4 (±1.6)	0.011 (±0)
	465	247 (±3)	3.5	70.5 (±1)	0.014 (±0)
<i>Aeromonas salmonicida</i>	405	345 (±131)	3.5	98.7 (±37)	0.011 (±0.001)
	465	555 (±10)	3.3	168.1 (±40.6)	0.006 (±0)
<i>Vibrio harveyi</i>	405	543.7 (±4.4)	4.3	126.4 (±1.0)	0.008 (±0)
	465	NA	NA	NA	NA
<i>Streptococcus parauberis</i>	405	446 (±66)	2.9	153.8 (±22.8)	0.007 (±0.001)
	465	743 (±153)	2.9	256.2 (±52.7)	0.004 (±0.001)

^aBactericidal efficiency, calculated as log₁₀ (10/N₀) per radiant flux (J cm⁻²)
 N₀ initial bacterial population in CFU ml⁻¹, NA not applicable, BE bactericidal efficiency

demonstrated that resistance plasmid for some antibiotics can be shared between bacterial fish pathogens, aquatic bacteria, and human pathogens, and some of them appear to have originated in the aquatic environment (Cabello et al. 2013). Thus, non-antibiotic therapies for infectious diseases are receiving considerable attention (Jori et al. 2006; Maisch 2009). It was previously demonstrated that blue light has a broad-spectrum bactericidal effect on both

Gram-negative and Gram-positive bacteria (Dai et al. 2012; Maclean et al. 2009). In this study, growth of the bacterial fish and shellfish pathogens evaluated was clearly inactivated by exposure to either a 405- or 465-nm LED light. Inactivation was dependent on light intensity and exposure time. Overall, our results show that Gram-negative bacteria, such as *P damsela* (36.1 J cm⁻²), *V. anguillarum* (41.2 J cm⁻²), and *E. tarda* (68.4 J cm⁻²),

Table 3 Relative growth of pathogenic bacteria with different initial population densities when exposed to 405- and 465-nm LED arrays for 24 h

Bacteria	Wavelength (nm)	Relative growth (%) ^a (± standard deviation) Initial bacterial cell counts (log CFU ml ⁻¹)					
		3	4	5	6	7	8
<i>Vibrio harveyi</i>	405	0 (±0) ^b	2 (±1) ^b	74 (±4) ^c	77 (±3) ^c	86 (±12) ^c	83 (±2) ^c
	465	70 (±61) ^b	68 (±2) ^b	72 (±2) ^b	69 (±3) ^b	78 (±18) ^b	76 (±2) ^b
<i>Vibrio anguillarum</i>	405	0 (±0) ^b	1 (±2) ^b	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	31 (±10) ^c
	465	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	107 (±40) ^c	107 (±30) ^c	105 (±25) ^c
<i>Edwardsiella tarda</i>	405	0 (±1) ^b	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	1 (±1) ^b	71 (±10) ^c
	465	1 (±1) ^b	0 (±0) ^b	76 (±18) ^c	121 (±24) ^d	86 (±6) ^c	83 (±2) ^c
<i>Aeromonas salmonicida</i>	405	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b
	465	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	44 (±4) ^c
<i>Photobacterium damsela</i>	405	77 (±3) ^b	105 (±2) ^d	109 (±12) ^d	103 (±4) ^d	86 (±8) ^{b, c}	99 (±1) ^{c, d}
	465	103 (±1) ^b	108 (±17) ^b	95 (±8) ^b	86 (±7) ^c	66 (±6) ^{b, c}	88 (±3) ^{b, c}
<i>Streptococcus iniae</i>	405	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	16 (±6) ^b	84 (±2) ^b	95 (±3) ^c
	465	1 (±1) ^b	0 (±0) ^b	0 (±0) ^b	1 (±0) ^c	2 (±1) ^c	51 (±8) ^d
<i>Streptococcus parauberis</i>	405	0 (±0) ^b	0 (±0) ^b	0 (±0) ^b	7 (±6) ^{b, c}	10 (±2) ^c	44 (±5) ^d
	465	0 (±0) ^b	0 (±1) ^b	0 (±0) ^b	11 (±2) ^c	10 (±4) ^c	44 (±6) ^d

Different letters (b, c, and d) in the superscripts indicate significant differences (P<0.05)
^aRatio of optical density (OD) of control to OD of LED treatment group (treatment OD_{630 nm}/control OD_{630 nm})

seem to be more sensitive to a 405-nm light than are Gram-positive bacteria like *S. parauberis* (153.8 J cm^{-2}) and *S. iniae* (90.4 J cm^{-2}). This result does not agree with a previous study which showed that Gram-positive bacteria such as *Staphylococcus*, *Clostridium*, and *Streptococcus* species were more susceptible to LED light than Gram-negative bacteria. Exceptions have been reported; *Enterococcus faecalis* suspensions exposed to 10 mW cm^{-2} light for up to 120 min experienced negligible inactivation (Maclean et al. 2009). Another study also found that the Gram-positive *Listeria monocytogenes* was more resistant to a 405-nm light than was the Gram-negative *Salmonella enterica* on acrylic and PVC surfaces (Murdoch et al. 2012). Taken together, it appears that Gram-positive bacteria are not always more rapidly inactivated than Gram-negative bacteria. The BE observed in this study are much lower than those seen in a previous study. This is because it took bacterial counts nine times over 200 min, which was much more frequent than in our study, where sampling was done only seven times over 48 h (Maclean et al. 2009). In addition, we used $250 \mu \text{ mol m}^{-2} \text{ s}^{-1}$ (approximately 7.4 mW cm^{-2}) and $516 \mu \text{ mol m}^{-2} \text{ s}^{-1}$ (approximately 13.3 mW cm^{-2}) intensities of 405- and 465-nm light, respectively, which are approximately 1.5–10 times lower than those used in previous studies (e.g., 19.5 mW cm^{-2} of 415 nm, 100 mW cm^{-2} of 415 or 455 nm, or 10 mW cm^{-2} of 405 nm) (Dai et al. 2013; Lipovsky et al. 2010; Maclean et al. 2009). This is one likely explanation as to why inactivation of pathogens in this study took longer than in previous studies. The precise mode of action of the antimicrobial effect of blue light is not yet fully understood. The commonly accepted hypothesis is that blue light excites endogenous intracellular porphyrins, which then behave as photosensitizers; photon absorption leads to energy transfer and, ultimately, the production of highly toxic reactive oxygen species (ROS) (Ashkenazi et al. 2003; Hamblin et al. 2005; Maclean et al. 2008). The differences in inactivation kinetics found in this study may be caused by organism-specific differences in porphyrin levels or porphyrin types, as suggested previously. The peak absorption wavelengths of different bacterial porphyrins may differ, and varying wavelengths may be required for their maximum photostimulation (Maclean et al. 2010). The degree of inactivation of bacterial suspensions with different initial densities was determined in order to assess LED activity on pathogens in the presence of nutrients mimicking a natural aquatic environment. *P. damsela* was able to survive a 405- or 465-nm light exposure when cultured on nutrient-enriched environment but was much more susceptible when suspended in PBS, as shown in Fig. 2. Several studies have reported that bacterial pathogens, including *Escherichia coli*, *A. salmonicida*, *Streptococcus pneumoniae*, and *V. harveyi*, produce different superoxide dismutase (SOD) and catalase isozymes inducible under certain growth conditions (Barnes et al.

1996; Flint et al. 1993; Vattanaviboon and Mongkolsuk 2001; Yesilkaya et al. 2000). However, *P. damsela* is not able to produce different SOD or catalase isozymes when exposed to oxidative stress induced by hydrogen peroxide, or under iron-depleted conditions (Díaz-Rosales et al. 2006). Also, *P. damsela*, possessing a high-affinity iron uptake system, grown under iron-limited conditions have a reduced amount of capsular material covering the cells (Do Vale et al. 2001; Naka et al. 2005). These indicate that *P. damsela* grown under nutrient-enriched conditions would be more resistant to oxidative stress (ROS) induced by LED irradiation than when grown under iron-limiting conditions (e.g., PBS). As it has been already demonstrated that blue light has caused no or very little damage to teleost (Migaud et al. 2007; Villamizar et al. 2011), it might be an alternative method to treat and prevent bacterial diseases in fish farm.

Conclusions

To the best of our knowledge, this study is the first to demonstrate that blue light is capable of inactivating major aquatic pathogens without requiring any external photosensitizer. As it is generally accepted that blue light is much less harmful to animal cells than is UV irradiation, and caused little damage to teleost that have already been demonstrated in previous studies (Migaud et al. 2007; Villamizar et al. 2011), application of blue light might be alternative to the use of antibiotics in aquaculture and would also have safety benefits. We hope our results will inspire further experiments to explore practical applications of blue light to fish and shellfish.

Acknowledgements

This research was a part of the project titled "LED-Marine Technology Convergence R&D Center" funded by the Ministry of Oceans and Fisheries, Korea.

Funding

This study was funded by the Ministry of Oceans and Fisheries, Korea.

Availability of data and materials

Considering this data will be used for enrolling patent, it will not be shared.

Authors' contributions

HJ mainly conducted the acquisition, analysis, and interpretation of the data and drafted and revised the manuscript. AR and GS participated in the design and cooperation of this work including acquiring the bacterial density and OD data. DH contributed to the conception and design of this work, the interpretation of the data, and the drafting and revising of the manuscript. All authors approved the final manuscript.

Competing interests

Authors HJ, AR, GS and DH are listed as inventors on a related patent application to the Korean Intellectual Property Office entitled "Inactivation methods aquatic bacterial pathogens using blue LED light" (Patent application number: 10-2015-0087358). The datasets supporting the conclusions of this article are available for non-commercial purpose upon request to the authors.

Received: 25 February 2016 Accepted: 5 August 2016

Published online: 30 August 2016

References

- Arrojado C, Pereira C, Tomé JPC, Faustino MAF, Neves MGPMS, Tomé AC, Cavaleiro JAS, Cunha A, Calado R, Gomes NCM, Almeida A. Applicability of photodynamic antimicrobial chemotherapy as an alternative to inactivate fish pathogenic bacteria in aquaculture systems. *Photochem Photobiol Sci*. 2011;10:1691–700.
- Ashkenazi H, Malik Z, Harth Y, Nitzan Y. Eradication of *Propionibacterium acnes* by its endogenous porphyrins after illumination with high intensity blue light. *FEMS Immunol Med Microbiol*. 2003;35:17–24.
- Baek GW, Kim JH, Gomez DK, Park SC. Isolation and characterization of *Streptococcus* sp. diseased flounder (*Paralichthys olivaceus*) in Jeju Island. *J Vet Sci*. 2006;7:53–8.
- Barnes AC, Horne MT, Ellis AE. Effect of iron on expression of superoxide dismutase by *Aeromonas salmonicida* and associated resistance to superoxide anion. *FEMS Microbiol Lett*. 1996;142:19–26.
- Bostock J, McAndrew B, Richards R, Jauncey K, Telfer T, Lorenzen K, Little D, Ross L, Handisye N, Gattward I, Corner R. Aquaculture: global status and trends. *Philos T Roy Soc B*. 2010;365:2897–912.
- Cabello FC, Godfrey HP, Tomova A, Ivanova L, Dölz H, Millanao A, Buschmann AH. Antimicrobial use in aquaculture re-examined: its relevance to antimicrobial resistance and to animal and human health. *Environ Microbiol*. 2013;15:1917–42.
- Dai T, Gupta A, Murray CK, Vrahas MS, Tegos GP, Hamblin MR. Blue light for infectious disease: *Propionibacterium acnes*, *Helicobacter pylori*, and beyond? *Drug Resist Updat*. 2012;15:223–36.
- Dai T, Gupta A, Huang YY, Tin R, Murray CK, Vrahas MS, Sherwood ME, Tegos GT, Hamblin MR. Blue light rescues mice from potentially fatal *Pseudomonas aeruginosa* burn infection: efficacy, safety, and mechanism of action. *Antimicrob Agents Chemother*. 2013;57:1238–45.
- Defoirdt T, Sorgeloos P, Bossier P. Alternatives to antibiotics for the control of bacterial disease in aquaculture. *Curr Opin Microbiol*. 2011;14:251–8.
- Díaz-Rosales P, Chabrilón M, Arijó S, Martínez-Manzanares E, Moriño MA, Balebona MC. Superoxide dismutase and catalase activities in *Photobacterium damsela* ssp. *Piscicida*. *J Fish Dis*. 2006;29:355–64.
- Do Vale A, Ellis AE, Silva MT. Electron microscopic evidence that expression of capsular polysaccharide by *Photobacterium damsela* subsp. *piscicida* is dependent on iron availability and growth phase. *Dis Aquat Organ*. 2001;44:237–40.
- Flint DH, Tuminello JF, Emptage MH. The inactivation of Fe-S cluster containing hydrolases by superoxide. *J Biol Chem*. 1993;268:22369–76.
- Ghanizadeh KE, Khodabandeh S. Effects of ultraviolet radiation on skin structure and ultrastructure in Caspian Sea salmon, *Salmo trutta caspius*, during alevin stage. *Toxicol Environ Chem*. 2010;92:903–14.
- Guardabassi L, Dalsgaard A, Raffatellu M, Olsen JE. Increase in the prevalence of oxolinic acid resistant *Acinetobacter* spp. Observed in a stream receiving the effluent from a freshwater trout farm following the treatment with oxolinic acid-medicated feed. *Aquaculture*. 2000;188:205–18.
- Hamblin MR, Viveiros J, Yang C, Ahmadi A, Ganz RA, Tolkoff MJ. *Helicobacter pylori* accumulates photoactive porphyrins and is killed by visible light. *Antimicrob Agents Chemother*. 2005;4:2822–7.
- Han HJ, Kim DH, Lee DC, Kim SM, Park SI. Pathogenicity of *Edwardsiella tarda* to olive flounder, *Paralichthys olivaceus*. *J Fish Dis*. 2006;29:601–9.
- Han HJ, Kim DY, Kim WS, Kim CS, Jung SJ, Oh MJ, Kim DH. Atypical *Aeromonas salmonicida* infection in the black rockfish, *Sebastes schlegelii* Hilgendorf, in Korea. *J Fish Dis*. 2011;34:47–55.
- Jori G, Fabris C, Soncin M, Ferro S, Coppellotti O, Dei D, Fantetti L, Chiti G, Roncucci G. Photodynamic therapy in the treatment of microbial infections: basic principles and perspective applications. *Laser Surg Med*. 2006;38:468–81.
- Kaweewat K, Hofer R. Effect of UV-B radiation on goblet cells in the skin of different fish species. *J Photochem Photobiol B*. 1997;41:222–6.
- Kleinpenning MM, Smits T, Frunt MHA, Van Erp PEJ, Van de Kerkhof PCM, Gerritsen RMJP. Clinical and histological effects of blue light on normal skin. *Photodermatol Photo*. 2010;26:16–21.
- Lipovsky A, Nitzan Y, Gedanken A, Lubart R. Visible light-induced killing of bacteria as a function of wavelength: implication for wound healing. *Laser Surg Med*. 2010;42:467–72.
- Maclean M, Macgregor SJ, Anderson JG, Woolsey G. The role of oxygen in the visible-light inactivation of *Staphylococcus aureus*. *J Photochem Photobiol B*. 2008;92:180–4.
- Maclean M, Macgregor SJ, Anderson JG, Woolsey G. Inactivation of bacterial pathogens following exposure to light from a 405-nanometer light-emitting diode array. *Appl Environ Microbiol*. 2009;75:1932–7.
- Maclean M, Macgregor SJ, Anderson JG, Woolsey GA, Coia JE, Hamilton K, Taggart I, Watson SB, Thakker B, Gettinby G. Environmental decontamination of a hospital isolation room using high-intensity narrow-spectrum light. *J Hosp Infect*. 2010;76:247–51.
- Maisch T. A new strategy to destroy antibiotic resistant microorganisms: antimicrobial photodynamic treatment. *Mini-Rev Med Chem*. 2009;9:974–83.
- Mata AI, Gibello A, Casamayor A, Blanco MM, Domínguez L, Fernández-Garayzábal JF. Multiplex PCR assay for detection of bacterial pathogens associated with warm-water streptococcosis in fish. *Appl Environ Microbiol*. 2004;70:3183–7.
- Migaud H, Cowan M, Taylor J, Ferguson HW. The effect of spectral composition and light intensity on melatonin, stress and retinal damage in post-smolt Atlantic salmon, *Salmo salar*. *Aquaculture*. 2007;270:390–404.
- Milton DL, Toole RO, Horstedt P, Wolf-Watz H. Flagellin A is essential for the virulence of *Vibrio anguillarum*. *J Bacteriol*. 1996;178:1310–9.
- Murdoch LE, Maclean M, Endarko E, Macgregor SJ, Anderson JG. Bactericidal effects of 405 nm light exposure demonstrated by inactivation of *Escherichia*, *Salmonella*, *Shigella*, *Listeria*, and *Mycobacterium* species in liquid suspensions and on exposed surfaces. *Sci World J*. 2012;2012:1–8.
- Naka H, Hirono I, Aoki T. Molecular cloning and functional analysis of *Photobacterium damsela* subsp. *piscicida* heme receptor gene. *J Fish Dis*. 2005;28:81–8.
- Osorio CR, Toranzo AE, Romalde JL, Barja JL. Multiplex PCR assay for ureC and 16S rRNA genes clearly discriminates between both subspecies of *Photobacterium damsela*. *Dis Aquat Organ*. 2000;40:177–83.
- Pang L, Zhang XH, Zhong Y, Chen J, Li Y, Austin B. Identification of *Vibrio Harveyi* using PCR amplification of the ToxR gene. *Lett Appl Microbiol*. 2006;43:249–55.
- Pridgeon JW, Klesius PH. Major bacterial diseases in aquaculture and their vaccine development. In: Hemming D, editor. *Anim Sci Rev* 2012. UK: CAB International; 2013. p. 141–56.
- Romalde JL. *Photobacterium damsela* subsp. *Piscicida*: an integrated view of a bacterial fish pathogen. *Int Microbiol*. 2002;5:3–9.
- Schmidt AS, Bruun MS, Dalsgaard I, Pedersen K, Larsen JL. Occurrence of antimicrobial resistance in fish-pathogenic and environmental bacteria associated with four Danish rainbow trout farms. *Appl Environ Microbiol*. 2000;66:4908–15.
- Subasinghe RP, Bondad-Reantaso MG, McGladdery SE. Aquaculture development, health and wealth. In: Subasinghe RP, Bueno P, Phillips MJ, Hough C, McGladdery SE, Arthur JR, editors. In proceedings of the Conference on Aquaculture in the Third Millennium: 20-25 February 2000. Bangkok: NACA and FAO; 2001. p. 167–91.
- Vattanaviboon P, Mongkolsuk S. Unusual adaptive, cross protection responses and growth phase resistance against peroxide killing in a bacterial shrimp pathogen. *Vibrio Harveyi*. *FEMS Microbiol Lett*. 2001;200:111–6.
- Villamizar N, Blanco-Vives N, Migaud H, Davie A, Carboni S, Sánchez-Vázquez FJ. Effects of light during early larval development of some aquacultured teleosts: a review. *Aquaculture*. 2011;315:86–94.
- Weinstein MR, Litt M, Kertesz DA, Wyper P, Rose D, Coulter M, McGeer A, Facklam R, Ostach C. Invasive infections due to a fish pathogen, *Streptococcus iniae*. *New Engl J Med*. 1997;28:589–94.
- Wklund T, Dalsgaard I. Occurrence and significance of atypical *Aeromonas salmonicida* in non-salmonid and salmoid fish species. *Dis Aquat Organ*. 1998;32:49–69.
- Won KM, Park SI. Pathogenicity of *Vibrio Harveyi* to culture marine fishes in Korea. *Aquaculture*. 2008;285:8–13.
- Yesilkaya H, Kadioglu A, Gingles N, Alexander JE, Mitchell TJ, Andrew PW. Role of manganese-containing superoxide dismutase in oxidative stress and virulence of *Streptococcus pneumoniae*. *Infect Immun*. 2000;68:2819–26.
- Yin Y, Gupta A, Hamblin MR. Light based anti-infectives: ultraviolet C irradiation, photodynamic therapy, blue light, and beyond. *Curr Opin Pharmacol*. 2013; 13:731–62.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

